

Detecting Classical Phase Transitions with Renyi Mutual Information

Jason Iaconis,^{1,2} Stephen Inglis,¹ Ann B. Kallin,¹ and Roger G. Melko^{1,3}

¹*Department of Physics and Astronomy, University of Waterloo, Ontario, N2L 3G1, Canada*

²*Department of Physics, University of California, Santa Barbara, CA, 93106-9530*

³*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada*
(Dated: October 31, 2012)

By developing a method to represent the Renyi entropies via a replica-trick on classical statistical mechanical systems, we introduce a procedure to calculate the Renyi Mutual Information in any Monte Carlo simulation. Through simulations on several classical models, we demonstrate that the Renyi Mutual Information can detect finite-temperature critical points, and even identify their universality class, without knowledge of an order parameter or other thermodynamic estimators. Remarkably, in addition to critical points mediated by symmetry breaking, the Renyi Mutual Information is able to detect topological vortex-unbinding transitions, as we explicitly demonstrate on simulations of the XY model.

Introduction – The universality and importance of the concept of information is exploited widely in mathematics and the physical sciences. Information sets the fundamental limits in communication (or uncertainty), regardless of system, technology, or physical material. Shannon was the first to quantify information using the concept of *entropy* – a quantity that has its roots in thermodynamics [1]. It is therefore not surprising that deep ties exist between the measurement of thermodynamic quantities and concepts associated with information theory.

There is already a rich cross-fertilization between ideas in condensed matter physics and the information sciences. Most recently, information measures have been used to quantify “hidden” correlations in materials – exchanges of information that can occur between two parts of a system that are *not* manifest in traditional condensed-matter estimators (such as correlation functions) [2]. For example, in spin liquid phases, correlation functions can rapidly decay as a function of spatial separation; however, due to constraints, hidden correlations exist across vast distances of the sample [3]. These can be manifest in entropy quantities measuring the amount of communication between two regions of the sample – resulting in a practically useful estimator for, among other things, topological order in condensed matter systems [4, 5].

Phase transitions offer another testing ground for the use of information quantities in condensed matter systems. Critical points are associated with a diverging correlation length, suggesting the existence of long-range channels for information transfer. However, it is not obvious that useful, measurable, quantities associated with this information can be exploited to tell us anything of practical importance about these phase transitions.

In this paper, we examine the Renyi Mutual Information (RMI), a measure that quantifies the amount of information contained in some region of a classical statistical mechanical system, about the rest of the system. Numerical measurements of classical mutual information [6, 7] typically calculate the reduced density matrix ex-

plicitly, or use symmetries to reduce its size, which give the von Neumann entropy directly but are computationally quite expensive. The RMI on the other hand is easily measured in standard Monte Carlo routines, requiring calculation via a *replica trick* [8, 9] rather than a reduced density matrix, making it immediately amenable to measurement on a vast number of models of interest to condensed matter, biophysics, and physical chemistry. We show that this RMI can be used in a practical way to identify finite-temperature phase transitions through finite-size scaling analysis on lattices of different sizes, *without* knowledge of an order parameter or any other thermodynamic quantity. The standard two-dimensional Ising model is used as a test case, demonstrating universality when results are compared to vastly different (even quantum mechanical [10]) models that exhibit the same universality class. Finally, we establish the ability of the method to detect the Berezinskii-Kosterlitz-Thouless transition, without relying on knowledge of any thermodynamic estimator such as the spin stiffness. This suggests the power of the RMI for detecting hidden transitions in a variety of other statistical mechanical models in the future.

Information, entropy, and the replica trick – Given a random variable X one can quantify its associated uncertainty, or equivalently, the amount of information one is missing by not knowing the state of X . There are various ways this can be measured, one of which is famously called *entropy*, embodied for example in the generalized Renyi entropies [11],

$$S_\alpha(X) = \frac{1}{1-\alpha} \ln \left(\sum_{i \in X} p_i^\alpha \right), \quad (1)$$

where p_i is the probability of outcome $X = i$. Taking the limit $\alpha \rightarrow 1$, one recovers Shannon’s familiar entropy, $S_1 = -\sum_i p_i \ln(p_i)$, which can be related to the thermodynamic entropy of a statistical mechanical system $S = \ln \Omega$, where Ω is the number of microstates, assuming all occur with equal probability. This relationship

is often exploited in the study of real physical systems through microscopic statistical mechanical models.

In such systems, one may also ask how much knowledge of a subsystem (call it A) is possible, assuming complete knowledge of another subsystem B . Although seemingly unmotivated in this language, correlation functions are a common example that incompletely quantify this knowledge. For the purposes of this paper, let us define a spatial subregion A as the complement of B , so that $A \cup B$ is the complete system; we could say that p_{i_A} is the probability of state i_A occurring in region A . The state i_A could be any classical (or indeed quantum) state: we will restrict our focus mainly to classical spin systems.

From Boltzmann, the probability of a given state in region A and region B for a classical spin system can be written

$$p_{i_A, i_B} = \frac{e^{-\beta E(i_A, i_B)}}{Z}, \quad (2)$$

where $E(i_A, i_B)$ is the energy of a system in states i_A in region A and i_B in region B , and $Z = \sum_{i_A, i_B} e^{-\beta E(i_A, i_B)}$ is the partition function. Eq. (2) is the quantity required to measure the entropy of the *entire* system – regions A and B . However, to obtain only the probability of a certain state in region A we can instead trace over all possible states in region B ,

$$p_{i_A} = \frac{\sum_{i_B} e^{-\beta E(i_A, i_B)}}{Z}. \quad (3)$$

Summing over the quantities in Eq. (2) or (3) raised to the power α , the Renyi entropies can be obtained in a straightforward way. In equilibrium statistical mechanical systems, physical estimators based on sums such as this are commonly calculated using importance sampling techniques, i.e. Markov Chain Monte Carlo.

In this paper we restrict our discussion to the *second* Renyi entropy which requires sums of the probabilities squared in Eq. (2) and (3). For example, to get the entropy of region A we use,

$$p_{i_A}^2 = Z^{-2} \left(\sum_{i_B} e^{-\beta E(i_A, i_B)} \right) \left(\sum_{j_B} e^{-\beta E(i_A, j_B)} \right), \quad (4)$$

leading to a Renyi entropy of

$$S_2(A) = -\ln \left(Z^{-2} \sum_{i_A} \sum_{i_B} \sum_{j_B} e^{-\beta(E(i_A, i_B) + E(i_A, j_B))} \right), \quad (5)$$

which can be sampled via a Monte Carlo simulation using the procedure described in the next section.

Note that, while the Renyi entropies tell us about uncertainty in the full system or part of the system, they do not reveal the correlations or information *between* two regions of a system. For this we must introduce the Renyi




FIG. 1: (color online). The replica trick for calculating $S_2(A)$ in a one-dimensional systems with 10 spins. In region A (the left 5 spins), configurations are constrained to always be identical. In region B , configurations are sampled independently in the two replicas.

Mutual Information (RMI),

$$I_\alpha(A; B) = S_\alpha(A) + S_\alpha(B) - S_\alpha(A \cup B). \quad (6)$$

This measure defines in a precise way the information that a full knowledge of B gives us about A , or vice versa, since it is symmetric in the regions A and B . One can see that the three constituents of $I_\alpha(A; B)$ are the so-called generalized Renyi Entropies defined above, including the entropy of region A , of region B , and of the full system $A \cup B$.

Monte Carlo Algorithm – For any physical system, the calculation of the Renyi entropy of a region, A , can be accomplished by sampling the quantity in Eq. (4) in a Monte Carlo simulation with a modified simulation geometry. Identifying the two individual sums in Eq. (4) with two independent *replicas* of the simulation immediately leads to a practical method for measurement of S_2 in Eq. (5), where the states of the subsystem B are sampled independently via two replicas (the sums over i_B and j_B), while the spin state in region A is constrained to be identical between these two replicas (the sum over i_A).

A schematic representation of a modified simulation is shown in Fig. 1. The constraint of identical region A s demands that, to be accepted, a spin update in region A must update spins in both replicas. For a spin in region A away from the boundary, because each replica of the system is at a temperature $1/\beta$, the acceptance rate of such a move has the form $e^{-2\beta\Delta E}$, where ΔE is the cost of the update in just one of the layers. This factor of 2 may be interpreted as moves in region A away from the boundary occurring at half the temperature as a move away from the boundary in region B .

The above method lets us generate states according to the probability in Eq. 4, but an additional step is needed to calculate the Renyi entanglement entropy. One approach, used previously [12], is to integrate the energy estimator, starting from infinite temperature, down to the temperature of interest, obtaining the Renyi entropy from both a replicated simulation and an unreplicated simulation. This necessitates a schedule of simulations over a large range of temperatures to gather enough detailed knowledge of the energy to integrate it. Alternatively, one may use methods able to directly extract the




FIG. 2: (color online). Left: The RMI per boundary length (I_2/ℓ) as a function of temperature for the Ising model. Dashed lines indicate T_c and $2T_c$. Upper right: the first derivative of the RMI for each system size as a function of temperature, with a fit to a logarithmic divergence shown in Eq. (8). Lower right: the temperature of the lower crossings of the RMI for sizes L and $2L$ as a function of $1/L$. Dashed line indicates T_c . Finite size scaling gives us $T_c = 2.2683(17)$.

ratio of partition functions at a given temperature. This is done by constructing moves that are able to connect simulations with different region sizes in a way that satisfies detailed balance (for a quantum approach, see [13]). By calculating the acceptance rate of moves connecting partition functions of differing region sizes, we can calculate the ratio of partition functions which can be related to the trace of higher powers of the density matrix [12].

Results on models – Using conventional Monte Carlo simulations of several models, we demonstrate the use of the second RMI to detect finite-temperature phase transitions. In the following, we use $I_2(A; B)$ where A and B are complementary regions, each defined as an $L \times L/2$ cylinder embedded in the $L \times L$ torus.

The first model we examine is the classical Ising model on a two-dimensional square lattice

$$H = -J \sum_{\langle ij \rangle} S_i^z S_j^z, \quad (7)$$

where $S_i^z = \pm 1/2$. There exists a transition to an ordered phase at a temperature $T_c/J = 2/\log(1 + \sqrt{2}) \approx 2.269$ [14], which offers us the simplest test for the RMI. Figure 2 illustrates $I_2(A; B)/\ell$ where ℓ is the length of the boundary between the regions (in all cases hereafter, $\ell = 2L$). Close inspection indicates approximate crossings of the $I_2(A; B)/\ell$ curves at T_c and $2T_c$. Examining the crossings as we move to larger system sizes we see that they extrapolate to the transition temperature as a function of $1/L$, as illustrated in the inset of Figure 2.

Remarkably, this behavior mimics the crossing seen previously in the analogous RMI quantity based on the *entanglement* entropies in a quantum spin-1/2 XXZ




FIG. 3: (color online). The RMI per boundary length (I_2/ℓ) as a function of temperature for the classical XY model, dashed lines indicate T_{BKT} and $2T_{BKT}$. Inset: The same plot for the quantum spin-1/2 XY model generated using a Wang-Landau technique [15]. Note that classically we reach much larger systems (only smallest and largest sizes are labeled).

model [12, 15]. That model also realizes a finite-temperature critical point, in the 2D Ising universality class (however at a different T_c , which is non-universal). There, it has been determined that the crossings were a manifestation of criticality, with scaling form,

$$I_2(A; B) = [c_1(t) + t \log t] \cdot \ell + c_2(t) + \mathcal{O}(1/\ell), \quad (8)$$

with $t = |T - T_c|/T_c$ [10]. It is important to note that $c_1(t)$ can be polynomial in t , where the strict “area law” at $t = 0$ is caused entirely by its constant piece, while divergences in derivatives of $I_2(A; B)$ are caused only by the $t \log t$ contribution. This diverging term is known from the $t \log t$ divergence in the boundary free energy as $t \rightarrow 0$ for the square lattice Ising model with a field applied to the boundary on an infinite half plane [16, 17]; we can thus use it as a test of universality in this model. In Figure 2, we plot the first derivative with respect to t of the RMI and see the formation of a $\log t$ singularity, in agreement with this scaling form. There, only the largest system studied has good agreement to this form – illustrating an advantage of using classical Monte Carlo over quantum Monte Carlo (QMC) techniques, the former being amenable to simulations on significantly larger system sizes.

We turn now to Monte Carlo simulations of the classical 2D XY model on a square lattice,

$$H = -J_{XY} \sum_{\langle ij \rangle} \cos(\theta_i - \theta_j). \quad (9)$$

This is a model with continuous spin variable $0 \leq \theta_i < 2\pi$ that undergoes a Berezinskii-Kosterlitz-Thouless (BKT) [18] transition from a phase with free vortices to one with bound vortex-antivortex pairs. Numerically, the detection of this phase transition is much more subtle

than the Ising phase transition, since the BKT transition has no local order parameter and is typically identified by the change from an exponential decay in correlations above the transition to a power law below [18]. Because the BKT transition exhibits this “essential singularity”, standard scaling theory does not work, making its detection via typical thermodynamic estimators (such as the specific heat) impossible. In 2D, this is circumvented by measuring the energy cost of uniformly twisting the spins in any lattice direction (the “spin stiffness”), and making use of a special universal “jump condition”, $T_{\text{BKT}} = \pi\rho_s/2$, due to Nelson and Kosterlitz [19]. By examining the spin stiffness in this way, the transition temperature has been found to be $T_{\text{BKT}} = 0.89294(8)$ [20]. However, one may wonder if the RMI can detect this phase transition without any need for specialized measurements like the above.

To address this, we examine the RMI for the classical XY model, shown in Figure 3. One can clearly see the development of crossings in the quantity I_2/ℓ near the value of T_{BKT} and $2T_{\text{BKT}}$. This is a strong indication that some sort of critical scaling is coming into play near the BKT transition. To examine this concept further, in the inset we illustrate the RMI for a completely different model, the quantum XY model $H = -t \sum_{\langle ij \rangle} (b_i^\dagger b_j + b_i b_j^\dagger)$, computed using Stochastic Series Expansion QMC where the quantum RMI is generated using a broad histogram approach [15]. In this case, similar crossings appear at the (non-universal) $T_{\text{BKT}} \approx 0.343$, giving strong evidence in support of the universality of this result that the RMI can detect the BTK transition.

However, unlike the relatively “clean” crossing of the 2D Ising critical point, these crossings have a larger finite-size scaling component. Figure 4 contains a detailed finite-size scaling analysis for the model in Eq. (9). This shows fits to two finite-size scaling forms which are derived by setting the correlation length near the critical temperature, $\xi = e^{-b/\sqrt{t}}(1 + \mathcal{O}(t))$ [21], equal to the linear system size L/L_0 with $t = (T - T_{\text{BKT}})/T_{\text{BKT}}$. Then by relating the temperature T with the finite-size transition temperature $T(L)$ gives:

$$T_{\text{BKT}}(L) = T_{\text{BKT}}(\infty) \left[1 + \frac{b^2}{\log^2(L/L_0)} + \frac{c}{\log^5(L/L_0)} \right] \quad (10)$$

Note, typical finite-size scaling analyses set the coefficient c equal to zero, effectively ignoring subleading $\mathcal{O}(t)$ corrections to the correlation length. Using this finite-size scaling from, the data for the crossing of I_2/ℓ convincingly approaches T_{BKT} in the limit of $L \rightarrow \infty$.

Discussion – The Renyi Mutual Information (RMI) is a quantity able to detect all correlations in a physical system, even those missed by traditional connected correlation functions. We have introduced a practical method to calculate the RMI using a modification of standard Monte Carlo techniques for classical statistical mechan-




FIG. 4: (color online). The crossing of the RMI for the classical XY model between sizes L and $2L$ as a function of $1/\log(L)^2$. Fit 1 fits the points to Eq. (10) assuming $c = 0$ and using the largest six systems while Fit 2 uses the full equation and all of the data. They give T_{BKT} estimate of $0.912(4)$ and $0.899(9)$, respectively. Inset: close-up of the lower crossing (only smallest and largest sizes are labeled). The arrows indicate the eight crossing values used in the main figure.

cal systems. We demonstrated that the RMI associated with the second Renyi entropy, I_2 , is able to identify both conventional critical points, as well as continuous transitions where standard scaling theory breaks down, namely the BKT transition. A straight-forward finite-size scaling analysis of I_2 is sufficient to identify each phase transition, without knowledge of an order parameter, broken symmetry, or critical theory.

The ease of implementation of the RMI measurement into any standard Monte Carlo routine could stimulate adoption to simulation studies of many fields of the physical sciences and beyond. Based on its ability to detect both Ising and BKT transitions, it would be interesting to implement the RMI measurement in simulations of generalized XY models where conventional estimators fail to identify such phase transitions [22]. The RMI will likely find great utility in other classes of classical models, such as loop [23, 24] or dimer models [25, 26], where universal properties of exotic criticality may be manifested [27]. Also particularly pressing is the question of whether RMI measurements can detect unconventional transitions in disordered systems, such as the glass [28, 29] or jamming transition [30]. Finally, the ubiquity of the Monte Carlo method in such far-reaching fields such as humanities or finance may precipitate the use of RMI in a host of unforeseen applications. In particular, for systems where analyzing mutual information in pre-existing data sets has proven illuminating, such as detecting transitions linked to financial market crashes [31], RMI may provide a *predictive* tool when used in conjunction with Monte Carlo simulations of models.

Acknowledgements – We thank R. Singh, P. Fendley, M. Müller, M. Gingras, S. Trebst, and A. Del Maestro for enlightening discussions. This work was made possible

by the computing facilities of SHARCNET. Support was provided by NSERC of Canada (A.B.K. and R.G.M.), the Ontario Ministry of Research and Innovation (J.I.), the Vanier Canada Graduate Scholarship program (S.I.), and the National Science Foundation under Grant No. NSF PHY11-25915 (R.G.M).

-
- [1] C. E. Shannon, The Bell System Technical Journal **27**, 379 (1948).
 - [2] M. M. Wolf, F. Verstraete, M. B. Hastings, and J. I. Cirac, Phys. Rev. Lett. **100**, 070502 (2008).
 - [3] X.-G. Wen, Int. J. Mod. Phys. B **4**, 239 (1990).
 - [4] M. Levin and X.-G. Wen, Phys. Rev. Lett **96**, 110405 (2006).
 - [5] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006).
 - [6] J. Wilms, J. Vidal, F. Verstraete, and S. Dusuel, Journal of Statistical Mechanics: Theory and Experiment **1**, 23 (2012).
 - [7] J. Wilms, M. Troyer, and F. Verstraete, Journal of Statistical Mechanics: Theory and Experiment **10**, 11 (2011).
 - [8] Y. Nakagawa, A. Nakamura, S. Motoki, and V. Zakharov (2009), arXiv:0911.2596.
 - [9] P. Buividovich and M. Polikarpov, Nuclear Physics B **802**, 458 (2008).
 - [10] R. R. P. Singh, M. B. Hastings, A. B. Kallin, and R. G. Melko, Phys. Rev. Lett. **106**, 135701 (2011).
 - [11] A. Renyi, Proc. of the 4th Berkeley Symposium on Mathematics, Statistics and Probability **1960**, 547 (1961).
 - [12] R. G. Melko, A. B. Kallin, and M. B. Hastings, Phys. Rev. B **82**, 100409 (2010).
 - [13] S. Humeniuk and T. Roscilde, ArXiv e-prints (2012), 1203.5752.
 - [14] L. Onsager, Phys. Rev. **65**, 117 (1944).
 - [15] S. Inglis and R. G. Melko, ArXiv e-prints (2012), 1207.5052.
 - [16] B. M. McCoy and T. T. Wu, Phys. Rev. **162**, 436 (1967).
 - [17] H. Au-Yang and M. E. Fisher, Phys. Rev. B **11**, 3469 (1975).
 - [18] J. M. Kosterlitz and D. J. Thouless, Journal of Physics C: Solid State Physics **6**, 1181 (1973).
 - [19] D. R. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).
 - [20] M. Hasenbusch, Journal of Physics A: Mathematical and General **38**, 5869 (2005).
 - [21] D. J. Amit, Y. Y. Goldschmidt, and S. Grinstein, Journal of Physics A: Mathematical and General **13**, 585 (1980).
 - [22] Y. Shi, A. Lamacraft, and P. Fendley, Phys. Rev. Lett. **107**, 240601 (2011).
 - [23] A. Nahum, J. T. Chalker, P. Serna, M. Ortuño, and A. M. Somoza, Phys. Rev. Lett. **107**, 110601 (2011).
 - [24] S. D. Geraedts and O. I. Motrunich, Phys. Rev. B **85**, 144303 (2012).
 - [25] D. Charrier, F. Alet, and P. Pujol, Phys. Rev. Lett. **101**, 167205 (2008).
 - [26] G. Chen, J. Gukelberger, S. Trebst, F. Alet, and L. Balents, Phys. Rev. B **80**, 045112 (2009).
 - [27] B. Swingle and T. Senthil (2011), arXiv:1109.3185.
 - [28] A. P. Young and H. G. Katzgraber, Phys. Rev. Lett. **93**, 207203 (2004).
 - [29] H. G. Katzgraber, D. Larson, and A. P. Young, Phys. Rev. Lett. **102**, 177205 (2009).
 - [30] A. Coniglio and M. Nicodemi, J. Phys.: Cond. Matt. **12**, 6601 (2000).
 - [31] M. Harre and T. Bossomaier, Europhysics Letters **87**, 18009 (2009).